

Upper bounds for the growth of Mordell-Weil ranks in pro- p towers of Jacobians

Jordan S. Ellenberg

January 24, 2010

Abstract

We study the variation of Mordell-Weil ranks in the Jacobians of curves in a pro- p tower over a fixed number field. In particular, we show that under mild conditions the Mordell-Weil rank of a Jacobian in the tower is bounded above by a constant multiple of its dimension. In the case of the tower of Fermat curves, we show that the constant can be taken arbitrarily close to 1. The main result is used in the forthcoming paper of Guillermo Mantilla-Soler on the Mordell-Weil rank of the modular Jacobian $J(Np^m)$.

1 Introduction

Let F be a number field, and let X/F be a smooth algebraic curve, not necessarily proper. Let p be a prime.

By a *pro- p tower* of curves over X we mean a diagram

$$\dots \rightarrow X_3/F \rightarrow X_2/F \rightarrow X_1/F \rightarrow X_0/F = X$$

where the map $X_n \rightarrow X$ is an étale cover, geometrically Galois with Galois group a finite p -group. Let T be a finite set of primes of $\mathcal{O}_F$ including p and ∞ . We say the tower *has good reduction* away from T if the diagram above extends to

$$\dots \rightarrow \mathcal{X}_3/R \rightarrow \mathcal{X}_2/R \rightarrow \mathcal{X}_1/R \rightarrow \mathcal{X}_0/R$$

where $R = \mathcal{O}_F[1/T]$, each $\mathcal{X}_n$ is smooth over R , and the maps are again finite étale.¹ Write G_F for the absolute Galois group of F and $G_T(F)$ for the Galois group of the maximal extension of F unramified outside T .

The main theorem of this paper is that the Mordell-Weil ranks of the Jacobians of the curves in a tower grow at most *linearly* in the genus.

Theorem (Theorem 3.1). *Write $\tilde{X}_n/F$ for the smooth proper curve over F containing X_n as open subscheme. Suppose G_F acts trivially on $H_1(X_{\bar{F}}, \mathbb{Z}/p\mathbb{Z})$. Then for any n and any subquotient A of the Jacobian of $\tilde{X}_n$ we have*

$$\text{rank } A(F) \leq 2 \dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{F}_p) \dim A.$$

¹A better definition might be to say that the tower has good reduction away from T if $X/\text{Spec } \mathcal{O}_F[1/T]$ is the complement in a proper smooth curve $\tilde{X}/\text{Spec } \mathcal{O}_F[1/T]$ of a divisor D which is also proper and smooth over $\text{Spec } \mathcal{O}_F[1/T]$; but the definition used here avoids technical issues orthogonal to the theme of the paper.

The upper bound for $\text{rank } A(F)/\dim A$ in Theorem 3.1 depends on X and p . In the special case of Fermat curves over $\mathbb{Q}$, we can do better.

Theorem (Theorem 4.1, special case $K = \mathbb{Q}$). *Let p be an odd prime, and let J_n be the Jacobian of the plane curve with equation*

$$x^{p^n} + y^{p^n} + z^{p^n} = 0$$

Then there is a constant C' , depending only on p , such that

$$\text{rank } J_n(\mathbb{Q}) \leq \dim J_n + C' p^n.$$

We note that $\dim J_n$ is on order p^{2n} .

1.1 Motivation and history

The original motivation for this paper comes from the theorem of Chabauty, and its improvement by Coleman [1], from which one obtains an upper bound for the number of points on $X(F)$ under the condition that the (generalized) Jacobian J of X satisfies the bound

$$\text{rank } J(F) < \dim J. \tag{1.1}$$

More recent work of Bruin, Flynn, Wetherell, and others has made it clear that Chabauty's method can be applied even when the inequality (1.1) fails for X , if it can be shown that it is satisfied for some set of étale covers of X . This leads naturally to the question

Question A: Does every curve X/F admit an étale cover Y/F such that $\text{rank } J(Y)(F) < \dim J(Y)$?

Any version of our main theorem giving a bound

$$\text{rank } A(F) \leq C \dim A + o(\dim A)$$

with $C < 1$ would imply an affirmative answer to Question A. Unfortunately, the constant we achieve is always at least 1; in the most favorable case, that of Fermat curves, we obtain a constant of 1, exactly at the Chabauty boundary.

The idea of this paper might be seen as an attempt at a “non-abelian Chabauty” method. For simplicity, we explain this in the case where $X = \mathbb{P}^1 - \{0, 1, \infty\}$ and $\{X_n\}$ is the tower of Fermat curves. Instead of studying the Selmer groups of the J_n individually, one might try to work “at the top” by studying a direct limit in m, n of $H^1(G_T(\mathbb{Q}(\zeta_{p^m})), J_n[p^\infty])$. This limit will be a cofinitely generated module for the non-abelian Iwasawa algebra $\Lambda = \mathbb{Z}_p[[\Gamma]]$, where $\Gamma = H_1(X_F, \mathbb{Z}_p) \rtimes \text{Gal}(\mathbb{Q}(\zeta_{p^\infty})/\mathbb{Q})$. One might hope that a “Selmer module” of this kind could be shown to be small in an appropriate sense, leading to an affirmative answer to Question A. The difficulty, of course, is that we have not placed any conditions on the restriction of our cohomology classes to decomposition groups at p . It is not clear to us exactly what conditions might be appropriate.

None of the Iwasawa-theoretic machinery is used in the present paper, but the argument should nonetheless be thought of as a finite-level approximation to the approach sketched above.

The main text of the present paper was written in 2002, at which time the author gave several lectures about the material contained here; we apologize for the long delay in making it publicly available. The intervening period saw the appearance of the beautiful work of Minhyong Kim [3], which can also be seen as a kind of non-abelian Chabauty – in his work, the quotient of the geometric étale fundamental group π by the m th term of its lower central series takes on the

role played by $\pi^{ab} = H_1(X_{\bar{F}}, \mathbb{Z}_p)$ in the classical method, and by $\pi/[[\pi, \pi], [\pi, \pi]]$ in the above paragraph. Under widely believed conjectures on Galois representations (and unconditionally in the case $X = \mathbb{P}^1 - \{0, 1, \infty\}$) Kim can show that the “unipotent analogue” of (1.1) is satisfied, and the finiteness of $X(F)$ follows.

The impetus for releasing the paper now is the recent work of Guillermo Mantilla-Soler [4], which uses Theorem 3.1 as an ingredient in an upper bound for the Mordell-Weil rank of the modular Jacobian $J(Np^m)$, as m grows with N fixed. The work of Mantilla-Soler combines the methods of the present paper with substantially more difficult group theory arising from the non-abelianness of the covers $X(Np^m)/X(N)$. It would be very interesting to revisit his work from the Iwasawa-theoretic viewpoint described above.

1.2 Acknowledgments

We are very grateful to Nigel Boston, John Coates, Guillermo Mantilla-Soler, William McCallum, and Pavlos Tzermias for useful discussions about the material in this paper. The author was partially supported by NSF-CAREER Grant DMS-0448750 and a Sloan Research Fellowship.

2 Unipotent Galois actions on fundamental groups

Let X/F be defined as in the previous section, and write π for the geometric étale fundamental group $\pi_1^{\text{ét}}(X/\bar{F})$. Let T be a finite set of primes including p, ∞ , and suppose

$$\dots \rightarrow X_3/F \rightarrow X_2/F \rightarrow X_1/F \rightarrow X_0/F = X$$

is a pro- p tower with good reduction away from T .

Write Θ^n for the Galois group of $(X_n)_{\bar{F}}/X_{\bar{F}}$, and Θ for the inverse limit of the Θ^n . Then Θ is a pro- p group admitting a surjection

$$\phi : \pi \rightarrow \Theta$$

and an action of G_F , arising from the compatible actions of G_F on the Θ^n .

Furthermore, Θ admits a filtration by finite-index normal subgroups

$$\Theta = \Theta_0 \supset \Theta_1 \supset \Theta_2 \supset \dots$$

where Θ_n is the kernel of the projection from Θ to Θ^n . Write H_n for $\phi^{-1}(\Theta_n)$. Then $H_n = \pi_1^{\text{ét}}(X_n/\bar{F})$.

Let V_n be the group H_n^{ab}/pH_n^{ab} . Then V_n is a finite-dimensional vector space over $\mathbb{F}_p$. Note that the action of G_F on V_n factors through $G_T(F)$, by the good reduction hypothesis.

Proposition 2.1. *Let V be a subquotient of V_n . Then*

$$\dim_{\mathbb{F}_p} H^1(G_T(F), V) \leq (\dim_{\mathbb{F}_p} V)(\dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{F}_p)).$$

Proof. We begin with a group-theoretic lemma.

Lemma 2.2. *The action of G_F on V_n is unipotent.*

Proof. The image of G_F on π lies in the kernel of

$$\mathrm{Aut}(\pi) \rightarrow \mathrm{Aut}(\pi^{ab}/p\pi^{ab})$$

which is a pro- p group. (See, e.g., [2, Th 12.2.2].) So the same is true for the image of G_F in $GL(V_n)$. But an action of a p -group on a finite-dimensional vector space over $\mathbb{F}_p$ is automatically unipotent. $\square$

By Lemma 2.2, we have a filtration of G_F -modules (whence also $G_T(F)$ -modules)

$$V_n = V_{n;0} \supset V_{n;1} \supset \dots V_{n;N} = 0.$$

whose successive terms yield exact sequences

$$0 \rightarrow V_{n;i+1} \rightarrow V_{n;i} \rightarrow (\mathbb{Z}/p\mathbb{Z})^{r_i} \rightarrow 0.$$

We thus obtain a cohomology sequence

$$H^1(G_T(F), V_{n;i+1}) \rightarrow H^1(G_T(F), V_{n;i}) \rightarrow H^1(G_T(F), (\mathbb{Z}/p\mathbb{Z})^{r_i}).$$

It follows that

$$\dim_{\mathbb{F}_p} H^1(G_T(F), V_{n;i}) \leq \dim_{\mathbb{F}_p} H^1(G_T(F), V_{n;i+1}) + r_i \dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{Z}/p\mathbb{Z})$$

and, by induction,

$$\dim_{\mathbb{F}_p} H^1(G_T(F), V_{n;i}) \leq \left(\sum_{j \geq i} r_j \right) \dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{Z}/p\mathbb{Z}).$$

The proposition follows by setting $i = 0$. $\square$

3 Towers of curves

We now explain how Proposition 2.1 can be used to give upper bounds for Mordell-Weil ranks (more precisely, Selmer ranks) in towers of curves. Let

$$\dots \rightarrow X_3/F \rightarrow X_2/F \rightarrow X_1/F \rightarrow X_0/F = X$$

be a pro- p tower with good reduction away from T , and define π, Θ_n, H_n, V_n as in the previous section.

Theorem 3.1. *Let $\tilde{X}_n/F$ be a smooth proper curve over F containing X_n as open subscheme, let J_n be the Jacobian of $\tilde{X}_n$, and let A be a subquotient of J_n . Suppose that G_F acts trivially on $H_1(X, \mathbb{F}_p)$. Then*

$$\frac{\mathrm{rank} A(F)}{\dim A} \leq 2 \dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{F}_p).$$

Proof. The usual descent on abelian varieties shows that

$$\mathrm{rank} A(F) \leq \dim_{\mathbb{F}_p} A(F)/pA(F) \leq \dim_{\mathbb{F}_p} H^1(G_T(F), A[p]).$$

The $G_T(F)$ -module $J_n[p]$ parametrizes étale abelian covers of $\tilde{X}_n$ with exponent p ; these restrict to étale abelian covers of X_n with exponent p . It follows that $J_n[p]$ is a quotient of V_n , whence $A[p]$ is a subquotient of V_n . The result now follows immediately from Proposition 2.1. $\square$

We give some examples where Theorem 3.1 applies.

Example 3.2. Let X/F be a proper curve of genus $g \geq 2$, admitting a smooth model over $\text{Spec } \mathcal{O}_F[1/T]$, and suppose $P \in X(F)$ is a rational point. Define X_n to be the Cartesian product

$$\begin{array}{ccc} X_n & \longrightarrow & J(X) \\ \downarrow & & \downarrow [p^n] \\ X & \longrightarrow & J(X) \end{array}$$

where the morphism $X \rightarrow J(X)$ is the Abel-Jacobi map sending Q to $[Q - P]$, and $[p^n]$ is multiplication by p^n . Then the X_n form a pro- p tower with good reduction away from T . In this case, $\Theta = \pi^{ab} \cong \mathbb{Z}_p^{2g}$. The condition that X is proper can be removed; see example 3.4 below for the simplest case.

Example 3.3. Let $X = Y(p)$ be the moduli space of elliptic curves with full level p structure, and let $X_n = Y(p^{n+1})$. Then the X_n form a pro- p tower with good reduction away from $T = p\infty$. Here, $\Theta = \Gamma(p)/\pm 1$, where $\Gamma(p)$ is the level p congruence subgroup of $\text{SL}_2(\mathbb{Z}_p)$. This case is treated in detail by Mantilla-Soler in [4].

Example 3.4. Let X be $\mathbb{P}^1 - \{0, 1, \infty\}$, which we write as the complement of the coordinate axes in the projective curve $x + y + z = 0$. Let X_n be the complement of the coordinate axes in the Fermat curve

$$x^{p^n} + y^{p^n} + z^{p^n} = 0.$$

Then the natural projection $X_n \rightarrow X$ is geometrically Galois with group $(\mathbb{Z}/p^n\mathbb{Z})^2$, and the X_n form a pro- p tower with good reduction away from $T = p\infty$. In this case, $\Theta = \mathbb{Z}_p^2$. Theorem 3.1 thus tells us that

$$\text{rank } J_n(\mathbb{Q}(\zeta_p)) \leq 2 \dim_{\mathbb{F}_p} H^1(G_T(\mathbb{Q}(\zeta_p)), \mathbb{F}_p) \dim J_n.$$

The constant on the right-hand side depends critically on the arithmetic of p ; in particular, it is large when the cyclotomic extension $\mathbb{Q}(\zeta_p)$ has a large p -class group. In the following section we will explain how to improve the above bound to one with milder dependence on p .

4 Towers of Fermat curves

In this section we prove the following theorem.

Theorem 4.1. *Let p be an odd prime, and let K be a number field such that the cyclotomic extension $K(\zeta_{p^\infty})/K(\zeta_p)$ has μ -invariant 0. Let J_n be the Jacobian of the Fermat curve $x^{p^n} + y^{p^n} + z^{p^n} = 0$. Then there exists a constant C' , depending on K and p , such that*

$$\text{rank}_{\mathbb{Z}} J_n(K) \leq [K : \mathbb{Q}] \dim J_n + C' p^n.$$

Let K be a number field, and let $X/K, X_n/K$ be as in Example 3.4. Let $F = K(\zeta_p)$, and let T be the set of primes dividing $p\infty$. We are now in the situation of Theorem 3.1, which tells us that

$$\text{rank } J_n(F) \leq 2 \dim J_n \dim_{\mathbb{F}_p} H^1(G_T(F), \mathbb{F}_p).$$

In other words, we have shown that the rank of J_n over F is bounded by a constant multiple c of its dimension, where c depends on p . We now turn to the more delicate problem of showing that c can be chosen independently from p .

Let $F_n = F(\mu_{p^n})$. Since Theorem 3.1 applies over any field containing μ_p , we also have

$$\text{rank } J_n(F_n) \leq 2 \dim J_n \dim_{\mathbb{F}_p} H^1(G_T(F_n), \mathbb{F}_p).$$

We have an inflation-restriction exact sequence

$$0 \rightarrow H^1(\text{Gal}(F_\infty/F_n), \mathbb{F}_p) \rightarrow H^1(G_T(F_n), \mathbb{F}_p) \rightarrow H^1(G_T(F_\infty), \mathbb{F}_p)^{\text{Gal}(F_\infty/F_n)} \rightarrow H^2(\text{Gal}(F_\infty/F_n), \mathbb{F}_p)$$

in which the last term is 0, since $\text{Gal}(F_\infty/F_n) \cong \mathbb{Z}_p$ has cohomological dimension 1. So

$$\dim_{\mathbb{F}_p} H^1(G_T(F_n), \mathbb{F}_p) = 1 + \dim_{\mathbb{F}_p} H^1(G_T(F_\infty), \mathbb{F}_p)^{G(F_\infty/F_n)}. \quad (4.2)$$

Furthermore, the long exact sequence

$$0 = H^0(G_T(F_\infty), \mathbb{Q}_p/\mathbb{Z}_p)/p \rightarrow H^1(G_T(F_\infty), \mathbb{Z}/p\mathbb{Z}) \rightarrow H^1(G_T(F_\infty), \mathbb{Q}_p/\mathbb{Z}_p)[p] \rightarrow 0$$

implies that $H^1(G_T(F_\infty), \mathbb{F}_p) = H^1(G_T(F_\infty), \mathbb{Q}_p/\mathbb{Z}_p)[p]$.

Now the group $H^1(G_T(F_\infty), \mathbb{Q}_p/\mathbb{Z}_p)$ is a well-understood object of classical Iwasawa theory. It is a cofinitely generated $\mathbb{Z}_p[[G(F_\infty/F)]]$ -module pseudo-isomorphic to

$$M^{r_2(F)} \oplus T$$

where M is cofree and T is cotorsion, and $r_2(F)$ denotes the number of complex places of F . Suppose the cyclotomic extension F_∞/F has μ -invariant 0 (as is the case, for instance, whenever $F/\mathbb{Q}$ is abelian by the theorem of Ferrero and Washington.) Then it follows from [5, 11.3.16, 11.3.17] that $T[p]$ is a finite group. We assume that the μ -invariant of F_∞/F is 0 from now on.

Since F contains μ_p , it is totally complex, and $r_2(F) = (1/2)[F : \mathbb{Q}]$. It follows that $H^1(G_T(F_\infty), \mathbb{F}_p)$ is pseudo-isomorphic to $M[p]^{(1/2)[F:\mathbb{Q}]}$.

Combining this with (4.2) and the fact that

$$M[p]^{\text{Gal}(F_\infty/F_n)} = \mathbb{F}_p[\text{Gal}(F_n/F)]$$

we have

$$\dim_{\mathbb{F}_p} H^1(G_T(F_n), \mathbb{Z}/p\mathbb{Z}) \leq (1/2)([F_n : \mathbb{Q}] + C)$$

for some constant C independent of n .

Applying Theorem 3.1, we now have:

Proposition 4.2. *Let $F \supset \mathbb{Q}(\zeta_p)$ be a number field whose cyclotomic extension has μ -invariant 0, and let A/F be a subquotient of the Jacobian of the Fermat curve $x^{p^n} + y^{p^n} + z^{p^n} = 0$, and let $F_n = F(\mu_{p^n})$. Then*

$$\text{rank } A(F_n) \leq (\dim A)([F_n : \mathbb{Q}] + C)$$

where C is a constant independent of n .

Recall that we have denoted the Galois group of $X_n/\bar{K} \rightarrow X/\bar{K}$ by Θ^n , which is isomorphic to $(\mathbb{Z}/p^n\mathbb{Z})^2$. Write H for $\text{Gal}(F_n/K)$, and G for the semidirect product $\Theta^n \rtimes H$ where H is given

its natural action on $\Theta^n \subset \text{Aut}(X_n/\bar{K})$. Then $J_n(F_n)$ carries an action of G , and the vector space $W = J_n(F_n) \otimes_{\mathbb{Z}} \mathbb{C}$ is a complex representation of G . Moreover,

$$\text{rank } J_n(F) = \dim W^H.$$

Since G is a semidirect product of an abelian group Θ^n by a subgroup of $\text{Aut}(\Theta_n)$, its irreducible representations are easy to describe (see [6, 8.2].) To be precise: let χ_i be a character of Θ^n , and let $H_i \subset H$ be the subgroup fixing χ_i . Then χ_i naturally extends to a character α_i of $G_i = \Theta^n H_i$. Let ψ be a character of H_i . Then $\text{Ind}_{G_i}^G(\alpha_i \otimes \psi)$ is an irreducible representation of G , denoted $\rho_{i,\psi}$. The dimension of $\rho_{i,\psi}$ is $[H : H_i]$, and the $\rho_{i,\psi}$ comprise all irreducible representations of G .

Note that

$$\text{Res}_H \rho_{i,\psi} = \text{Res}_H \text{Ind}_{G_i}^G(\alpha_i \otimes \psi) = \text{Ind}_{H_i}^H \psi$$

by [6, 7.3]. So $\dim(\rho_{i,\psi})^H$ is 1 if ψ is trivial, and 0 otherwise. In any case,

$$\dim(\rho_{i,\psi})^H \leq ([H : H_i]^{-1}) \dim \rho_{i,\psi}. \quad (4.3)$$

Let B be the kernel of the map $\Theta^n \rightarrow \Theta^{n-1}$. Then we have an exact sequence of $\mathbb{C}[H]$ -modules

$$0 \rightarrow W^B \rightarrow W \rightarrow W' \rightarrow 0.$$

We note, first of all, that W^B is precisely $J_{n-1}(F_n) \otimes_{\mathbb{Z}} \mathbb{C}$. The map $X_n \rightarrow X_{n-1}$ induces a homomorphism $J_{n-1} \rightarrow J_n$, whose cokernel we call J'_n . We also write $J'_0 = J$. Then W' is isomorphic to $J'_n(F_n) \otimes_{\mathbb{Z}} \mathbb{C}$.

Now W' is the direct sum of all the irreducible constituents of W on which B does not act trivially. The $\rho_{i,\psi}$ on which B does not act trivially are precisely those for which χ_i has exact order p^n . Note that the stabilizer H_i is trivial for any such χ_i . Applying (4.3) to each constituent of W' , we find that

$$\dim(W')^H \leq |H|^{-1} \dim W' = [F_n : K]^{-1} \dim W'.$$

Combining this with the upper bound on $\dim W'$ given by Proposition 4.2, we have

$$\dim(J'_n(K) \otimes_{\mathbb{Z}} \mathbb{C}) = \dim(W')^H \leq ([K : \mathbb{Q}] + C[F_n : K]^{-1})(\dim J'_n).$$

Now J_n is isogenous to $\oplus_{i=0}^n J'_i$. It follows that

$$\text{rank } J_n(K) \leq [K : \mathbb{Q}] \dim J_n + C \sum_{i=0}^n [F_i : K]^{-1} \dim J'_i.$$

The sum over i on the right-hand side is bounded above by a constant multiple of p^n . This completes the proof of Theorem 4.1.

References

- [1] R.F. Coleman. Effective Chabauty. *Duke Math. J.*, 52(3):765–770, 1985.
- [2] M. Hall. *The theory of groups*. Chelsea Publishing Company, Incorporated, 1976.
- [3] Minhyong Kim. The unipotent Albanese map and Selmer varieties for curves. *Publ. RIMS*, 45(1):89–133, 2009.

- [4] Guillermo Mantilla-Soler. *Mordell-Weil ranks in towers of modular Jacobians*. PhD thesis, U. Wisconsin-Madison, 2010.
- [5] J. Neukirch, A. Schmidt, and K. Wingberg. *Cohomology of number fields*. Springer Verlag, 2008.
- [6] J.P. Serre. *Linear representations of finite groups*. Springer Verlag, 1977.